

dowagro.ca

For more information on crop groups, see the reference section of the Dow AgroSciences horticulture guide or call the Solutions Center at 1-800-667-3852.



Solutions for the Growing World

EXCELLENT HERBICIDES FOR FRUITS

Goal™ 2XL
EMULSIFIABLE CONCENTRATE HERBICIDE

- Group 14 herbicide
- Excellent pre- and post-emergence control of broadleaf weeds
- Pre-harvest interval: raspberries = 50 days; high bush blueberries = 50 days; strawberries = 150 days

Lontrel™ 360
HERBICIDE

Lontrel™ XC
HERBICIDE

- Group 4 herbicide
- The standard for thistle control in strawberries, blueberries and cranberries
- Translocates for complete in-season thistle control
- Pre-harvest interval: cranberries = 60 days; low bush blueberries = 10 months

DuPont™ Fontelis®
fungicide

- Group 7 fungicide
- Excellent residual, preventative and post-infection activity in many crops
- Provides translaminar and locally systemic protection
- Post-harvest interval: berries, blueberries, stone fruits = 0 days; pome fruits = 28 days

EXCELLENT FUNGICIDES FOR FRUITS

Quintec™
FUNGICIDE

- Group 13 fungicide
- Protectant powdery mildew control using a unique mode of action
- Preventative applications on strawberries can be applied every 10-14 days
- Pre-harvest interval: strawberries = 1 day

Nova™
FUNGICIDE

- Group 3 fungicide
- Systemic, providing long-lasting, effective powdery mildew control
- Can be re-applied in blueberries every 7-14 days; strawberries every 14-21 days
- Pre-harvest interval: strawberries = 3 days; bush berries = when harvesting by hand 6 days except for low bush blueberries, which is 4 days; when harvesting mechanically 1 day; cranberries = when harvesting by hand 6 days; when harvesting mechanically 1 day

DuPont™ Tanos®
fungicide

- Group 11 and Group 27 fungicide
- Post-infection and kick-back activity against late blight and control of late blight stem infections
- Offers both systemic and protectant activity
- Post-harvest interval: cane berries = 9 days



HIGH PERFORMANCE PRODUCTS

from Dow AgroSciences for fruits

Solutions for the Growing World

FRUIT GROWER PRODUCT GUIDE	INSECTICIDES										FUNGICIDES					HERBICIDES						
	Close [®]	Delegate [®]	Entrust [®]	GF-120 Fruit Fly Bait	Intrepid [®]	Lannate [®]	Lorsban [®] 50W	Lorsban [®] NT	Success [®]	TwinGuard [®]	Vydate	Dithane [®] Rainshield [™]	Fontelis [®]	Indar [®]	Nova [®]	Quintec [®]	Tanos [®]	Goal [®] 2XL	Kerb [®] SC	Lontrel [®] 360/Lontrel [®] XC	Prism [®] SG	VP480
Apple	x	x	x	x	x	x			x	x	x	x		x				x	x			x
Apricot	x	x	x		x				x	x			x	x	x	x				x		x
Blackberry		x	x		x				x						x		x					
Blueberry – lowbush		x	x	x	x				x				x		x		x	x	x			x
Blueberry – highbush		x	x	x	x				x					x	x				x	x	x	
Cantaloupe					x							x	x		x	x						
Cherry	x	x	x	x	x				x	x			x	x	x	x				x		x
Cranberry		x	x		x				x				x							x		x
Grape	x	x	x		x				x			x			x	x						x
Nectarine	x	x	x		x		x		x	x			x	x	x	x				x		
Peach	x	x	x		x		x		x	x			x	x	x	x				x		x
Pear	x	x	x		x				x	x			x		x				x			x
Plum/Prune	x	x	x		x				x	x			x	x		x				x		x
Raspberry		x	x		x				x		x				x		x	x				x
Saskatoon berry		x	x		x				x						x							
Strawberry		x	x				x	x	x				x		x	x		x	x	x		x
Watermelon					x							x	x		x	x						

*Non-bearing trees.

Always read the label for rates, instructions and precautions. Please call the Solutions Center at 1-800-667-3852 or visit dowagro.ca

Success[™]
INSECTICIDE

Delegate[™]
INSECTICIDE

Proven, excellent control of a broad range of pests

- Group 5 insecticide
- Excellent control of a broad range of insects including spotted wing drosophila
- Unique mode of action that controls insects two ways – on contact and by ingestion – for quick knockdown and residual activity
- Moves through the leaf, providing extra protection against insects that feed from the underside of leaves and increasing resistance to wash-off by rain
- Pre-harvest interval: strawberries, cane berries = 1 day; bush berries = 3 days; cranberries = 21 days

Entrust[™]
INSECTICIDE

Improved control of insects on organic farms

- Group 5 insecticide
- Excellent activity on armyworm, thrips, leafminers, codling moth, cherry fruit fly and spotted wing drosophila
- Provides quick knockdown and residual activity because it controls insects on contact and by ingestion
- Pre-harvest interval: low-growing berry subgroup and cane berry subgroup = 1 day; bush berries and low-growing berries = 3 days; cranberries: 21 days

Intrepid[™]
INSECTICIDE

Controls pests at early stages to get your berries off to a strong start

- Group 18 insecticide
- Induces premature molt in lepidopterous pests (caterpillars) without adversely affecting beneficial insects such as bees, predatory mites, bees, wasps and spiders
- Has both ovicidal and larvicidal activity providing long residual control (up to 10-14 days after application)
- Unique mode of action and valuable tool in any IPM program
- Pre-harvest interval: blueberries = 7 days; cranberries = 14 days